

References Cited

- AFWA (Association of Fish and Wildlife Agencies). 2007. Guidance for Wildlife Action Plan (Comprehensive Wildlife Conservation Strategy) review and revisions. Available from <http://www.teaming.com/sites/default/files/Revision%20Guidance%20Letter%20NAAT.pdf>
- AFWA. 2011. Measuring the effectiveness of State Wildlife Grants: final report. Available from http://www.fishwildlife.org/files/Effectiveness-Measures-Report_2011.pdf
- AFWA, Teaming With Wildlife Committee, State Wildlife Action Plan (SWAP) Best Practices Working Group. 2012. Best practices for State Wildlife Action Plans—voluntary guidance to States for revision and implementation. Washington (DC): Association of Fish and Wildlife Agencies 80 p. Available from <http://www.fishwildlife.org/files/SWAPBestPractices.pdf>
- Ahlstedt, S.A., J.R. Powell, R.S. Butler, M.T. Fagg, D.W. Hubbs, S.F. Novak, S.R. Palmer, and P.D. Johnson. 2004. Historical and current examination of freshwater mussels (Bivalvia: Margaritiferidae, Unionidae) in the Duck River basin Tennessee. Contract report for the Tennessee Wildlife Resources Agency FA-02-14725-00.
- Anderson, M.G., A. Barnett, M. Clark, C. Ferree, A. O. Sheldon, and J. Prince. 2014. Resilient sites for terrestrial conservation in the Southeast Region. The Nature Conservancy, Eastern Conservation Science 127 p. Available from https://easterndivision.s3.amazonaws.com/Terrestrial/Resilient_Sites_for_Terrestrial_Conservation_In_the_Southeast_Region.pdf
- Andrews, K.M., T. M. Norton, J. E. Colbert, S. E. Nelson. 2013. Survivorship, Home Range Size, and Health of Head-Started Juvenile Eastern Box Turtles (*Terrapene carolina*). Jekyll Island Georgia Sea Turtle Center.
- Arnwine, D H., K J. Sparks, and R R. James. 2006. Probabilistic monitoring of streams below small impoundments in Tennessee. Nashville (TN): Tennessee Department of Environment and Conservation. Available from http://www.tn.gov/assets/entities/environment/attachments/isp_report.pdf
- AWEA (American Wind Energy Association). 2015. Get the facts. [Consulted 29 May 2015]. Available from <http://www.awea.org/Resources/Content.aspx?ItemNumber=900&navItemNumber=587>
- Bailey, B. 1999. Social and economic impacts of wild harvested products [dissertation]. West Virginia University-Morgantown.
- Bailey, R.G., P.E. Avers, T. King, and W.H. McNab. 1994. Ecoregions and subregions of the United States (map). Washington, DC: U.S. Geological Survey. Scale 1:7,500,000. Colored. Accompanied by a supplementary table of map unit descriptions compiled by W.H. McNab, and R.G. Bailey, editors. Prepared for the USDA Forest Service.

Benbrook, C. 2009. Impacts of genetically engineered crops on pesticide use in the United States: the first thirteen years. The Organic Center 62 p. Available from <https://www.organic-center.org/reportfiles/GE13YearsReport.pdf>

Benbrook, C. 2012. Impacts of genetically engineered crops on pesticide use in the U.S. -- the first sixteen years. Environmental Sciences Europe 24:24. doi:10.1186/2190-4715-24-24. Available from <http://www.enveurope.com/content/24/1/24>

Bhatta, B. 2010. Causes and consequences of urban growth and sprawl. In: Bhatta, B, editor. Analysis of urban growth and sprawl from remote sensing data. Berlin Heidelberg: Springer-Verlag. p. 17-36.

BLM (Bureau of Land Management). Split estate. [Consulted 29 May 2015]. Available from http://www.blm.gov/wo/st/en/prog/energy/oil_and_gas/best_management_practices/split_estate.print.html

Bowen, K.D., P. L. Colbert, AND F.J. Janzen. 2004. Survival and Recruitment in a Human-Impacted Population of Ornate Box Turtles, *Terrapene ornata*, with Recommendations for Conservation and Management. J. of Herpetology, Vol. 38, No. 4, pp. 562-568, 2004

Bradley, M.W. and G.E. Hileman. 2006. Sinkhole flooding in Murfreesboro, Rutherford County, Tennessee, 2001-02. U.S. Geological Survey Scientific Investigations Report 2005-5281. Available from <http://pubs.usgs.gov/sir/2005/5281/PDF/SIR20055281.pdf>

Burger, L.W., Jr. 2006. Creating wildlife habitat through Federal Farm Programs: an objective-driven approach. Wildlife Society Bulletin 34: 994-999.

Butler, R.S. 2002. Imperiled fishes of the Southern Appalachian Ecosystem, with emphasis on the non-federally listed fauna. Asheville (NC): prepared for the Southern Appalachian Ecosystem Team, U.S. Fish and Wildlife Service.

Cox, R.E. 2014. Potential impacts of increased feral pig populations to Virginia's National Forests. Virginia Polytechnic Institute and State University. Available from <https://scholar.vt.edu/access/content/group/5b95dc6f-a3ef-4ce5-8e1a-875819148663/Web/Student%20Publications/RobertCox.Capstone2014.pdf>

Cumberlandian Region Mollusk Restoration Committee. 2010. Plan for the population restoration and conservation of freshwater mollusks of the Cumberlandian Region. V + 145 pp. Available from <http://applcc.org/projects/trb/resources/reports-and-reference-documents/plan-for-the-population-restoration-and-conservation-of-imperiled-freshwater-mollusks-of-the-cumberland-region/view>

Dahl, T.E. 1990. Wetlands losses in the United States, 1780's to 1980's. Washington (DC): U.S. Department of the Interior, Fish and Wildlife Service. Available from <https://www.fws.gov/wetlands/Documents/Wetlands-Losses-in-the-United-States-1780s-to-1980s.pdf>

Dale, V.H., K.L. Kline, J. Wiens, and J. Fargione. 2010. Biofuels: implications for land use and biodiversity. Biofuels and Sustainability Reports. Ecological Society of America. Available from http://web.ornl.gov/sci/ees/cbes/Publications/ESA%20Biofuels%20Report_VH%20Dale%20et%20al%202010.pdf

DOI (U.S. Department of the Interior), U.S. Fish and Wildlife Service, and U.S. Department of Commerce, U.S. Census Bureau. 2011. National survey of fishing, hunting, and wildlife-associated recreation. Revised February 2014. Available from <https://www.census.gov/prod/2012pubs/fhw11-nat.pdf>

DOD (U.S. Department of Defense). 2006. Arnold Air Force Base: natural resources conservation – large installation /civil works facility. Available from http://www.denix.osd.mil/awards/upload/NRC_Inst_Arnold.pdf

DOD Arnold Engineering Development Center. 2006. Integrated natural resources management plan. Tennessee (USA): Arnold Air Force Base.

Drohan P.J., M. Brittingham, J. Bishop, and K. Yoder. 2012. Early trends in landcover change and forest fragmentation due to shale-gas development in Pennsylvania: a potential outcome for the Northcentral Appalachians. Environmental Management 49(5):1061-1075.

EGCPJV (East Gulf Coastal Plain Joint Venture). 2014. A burning issue: prescribed fire and fire-adapted habitats of the East Gulf Coastal Plain. Available from <https://drive.google.com/file/d/0B7OEhGi--LTCQjVpcDltSnBQTzA/view>

Elliott, K. G. 2010. Celebrating the Tenth Anniversary of State Wildlife Grants in Tennessee: a new wildlife tradition of keeping “common species common” is key to the ecosystems and economic health of the Volunteer State. Tennessee Wildlife Resources Agency Special Report. Available from <http://www.state.tn.us/twra/cwcs/swg10years.pdf>

Entrekin, S., M. Evans-White, B. Johnson, and E. Hagenbuch. 2011. Rapid expansion of natural gas development poses a threat to surface waters. Frontiers in Ecology and the Environment 9(9):503-511. Available from <http://www.cce.cornell.edu/EnergyClimateChange/NaturalGasDev/Documents/PDFs/Entrekin%20et%20al%20Frontiers%20in%20Ecology%20and%20the%20Environment.pdf>

EPA (Environmental Protection Agency). 2002. Clinch and Powell Valley Watershed ecological risk assessment. Available from <http://cfpub.epa.gov/ncea/cfm/recorddisplay.cfm?deid=15219>

EPA. 2012. Acid rain. [Consulted 30 May 2015]. Available from <http://www.epa.gov/acidrain/>

EPA. 2014a. Basic information about Polychlorinated Biphenyls (PCBs) in drinking water. [Consulted 28 May 2015]. Available from <http://water.epa.gov/drink/contaminants/basicinformation/polychlorinated-biphenyls.cfm>

EPA. 2014b. Mercury. [Consulted 31 May 2015]. Available from <http://www.epa.gov/mercury/about.htm>

EPA. 2014c. Section 319 Nonpoint Source Program success story, Tennessee. Available from http://water.epa.gov/polwaste/nps/success319/upload/tn_craborchard-2.pdf

EPA. 2015. Watershed assessment, tracking & environmental results (WATERS): Tennessee water quality assessment report. [Consulted 8 July 2015]. Available from http://ofmpub.epa.gov/waters10/attains_state.control?p_state=TN

Etnier, D.A. and W.C. Starnes. 1993. The fishes of Tennessee. Knoxville (TN): University of Tennessee Press 681 p.

Ewing, R., J. Kostyack, D. Chen, B. Stein, and M. Ernst. 2005. Endangered by sprawl: how runaway development threatens America's wildlife. Washington (D.C.): National Wildlife Federation, Smart Growth America, and NatureServe. Available from <http://www.nwf.org/~media/PDFs/Global-Warming/Reports/EndangeredbySprawl.pdf>

Fiedler, J. 2004. Assessment of bat mortality and activity at Buffalo Mountain Windfarm, eastern Tennessee (thesis). The University of Tennessee-Knoxville. Available from http://www.tva.gov/environment/bmw_report/bat_mortality_bmw.pdf

Fisher, M. Fracking's footprint: scientists study impact of shale gas development on Pennsylvania's forests. 2012. CSA News. Available from <https://www.crops.org/files/publications/csa-news/frackings-footprint.pdf>

Glick, P., S.R. Palmer and J.P. Wisby. 2015. Climate Change Vulnerability Assessment for Tennessee Wildlife and Habitats. Report prepared by the National Wildlife Federation and The Nature Conservancy-Tennessee Chapter for the Tennessee Wildlife Resources Agency, Nashville, TN. Available from http://tnswap.com/files/TNSWAP_2015VulnerAssmt.pdf

Global Ranavirus Consortium. 2015. Introduction. [Consulted 26 July, 2015]. Available from <http://www.ranavirus.org>

Granstaff, E.M., J.S. Perkin, and T.Roberts. 2015. A GIS tool for prioritizing dams for removal within the Tennessee and Cumberland River Basins. Tennessee Technological University.

Grimm, L.T. 2002. Certifying hydropower for “green” energy markets: the development, implementation, and future of the low impact hydropower certification program. Low Impact Hydropower Institute. Available from <http://www.lowimpacthydro.org/assets/files/Program%20Documents/Lydia%20Grimm%20on%20LIHI%20formation%202002.pdf>

Gunnarsson, B. and J. Johnsson. 1989. Effects of simulated acid rain on growth rate of a spruce-living spider. *Environmental Pollution* 56(4):311-317.

Hannah, T., R. Colvin, and P. English. 2013. Climate change vulnerability assessment report for high priority wildlife species in West Tennessee. Available from https://docs.google.com/file/d/0B_L0agRG2cxYNTUtUUZicjNRbE0/edit

Harper, C. and J. Birckhead. 2012. Use of prescribed fire in Tennessee and the Tennessee Prescribed Burning Act. *Treeline* 9(2):7-9. Available from http://fwf.ag.utk.edu/personnel/charper/pdfs/TN_Rx_Fire_Law_TreeLine_June2012.pdf

Harvey, M.J. and E.R. Britzke. 2002. Distribution and status of endangered bats in Tennessee. Final Report Prepared for the Tennessee Wildlife Resources Agency 44 p.

Hutson, S.S., M.C. Koroa, and C.M. Murphree. 2004. Estimated use of water in the Tennessee River Watershed in 2000 and projections of water use to 2030. USGS Water-Resources Investigations Report 03-4302. Available from <http://pubs.usgs.gov/wri/wri034302/PDF.htm>

Jolley, D.B., S.S. Ditchkoff, B.D. Sparklin, L.B. Hanson, M.S. Mitchell, and J.B. Grand. 2010. Estimate of herpetofauna depredation by a population of wild pigs. *Journal of Mammalogy* 91(2):519-524.

Joyce, L.A., C.H. Flather, and M. Koopman. 2008. Final Project Report, WHPRP, 1.B: Analysis of Potential Impacts of Climate Change on Wildlife Habitats in the U.S. Available from http://ncseonline.org/sites/default/files/1B_ljoyce_WHPRP%20Final%20Report%20Project%20for%20web.pdf

Kershner, J., and E. Mielbrecht. 2012. A climate-informed conservation blueprint for the Greater Puget Sound Ecoregion. EcoAdapt and Geos Institute.

Keys, J. Jr., C. Carpenter, S. Hooks, F. Koenig, W.H. McNab, W. Russell, M.L. Smith. 1995. Ecological units of the eastern United States – first approximation (cd-rom), Atlanta (GA): U.S. Department of Agriculture, Forest Service. GIS coverage in ARCINFO format, selected imagery and map unit tables. Available from <http://www.srs.fs.usda.gov/econ/data/keys/readme.1st>

Lamb, J.W. and G.R. Wyckoff, Eds. 2010. Cooperative White-nose Syndrome Monitoring and Surveillance Plan for Tennessee. Available from <http://www.tnbgw.org/wnstnplan%202010.pdf>

Langwig, K.E., J. Voyles, M.Q. Wilber, W.F. Frick, K.A. Murray, B.M. Bolker, J.P. Collins, T.L. Cheng, M.C. Fisher, J.R. Hoyt, D.L. Lindner, H.I. McCallum, R. Puschendorf, E.B. Rosenblum, M. Toothman, C.K.R. Willis, C.J. Briggs, and A.M. Kilpatrick. 2015. Context-dependent conservation responses to emerging wildlife diseases. *Frontiers in Ecology and the Environment* 13:195-202. Available from <http://www.esajournals.org/journals.conserveonline.org:2048/doi/full/10.1890/140241>

Lannoo, M.J. 2005. *Amphibian declines: the conservation status of U.S. amphibians*. Berkeley (CA): University of California Press 1094 p.

Marion J.L. and N. Olive. 2006. *Assessing and Understanding Trail Degradation: Results from Big South Fork National River and Recreational Area*. USGS Patuxent Wildlife Research Center Cooperative Park Studies Unit. Virginia Technological University, Department of Forestry. Blacksburg, VA 84 p. Available from http://www.pwrc.usgs.gov/prodabs/pubpdfs/6612_marion.pdf

Master, L.L., S.R. Flack and B.A. Stein. 1998. *Rivers of life: critical watersheds for protecting freshwater biodiversity*. Arlington (VA): The Nature Conservancy.

McKinney, M.L. 2008. Effects of urbanization on species richness: a review of plants and animals. *Urban Ecologist* 11:161-176. Available from <http://www.mit.edu/people/spirn/Public/Granite%20Garden%20Research/Urban%20ecology/McKinney%202008%20Species%20Richness.pdf>

Medlock, K., S. R. Palmer and J.P. Wisby. 2013. *Tennessee Freshwater Mollusk Strategic Plan*. The Nature Conservancy, Nashville, TN. 44 pp.

Melillo, J.M., T. (T.C.) Richmond, and G.W. Yohe, editors. 2014: *Climate change impacts in the United States: the Third National Climate Assessment*. U.S. Global Change Research Program 841 p. doi:10.7930/J0Z31WJ2. Available from <http://nca2014.globalchange.gov/downloads>

Mitchell, J.C., T.K. Pauley, D.I. Withers, S.M. Roble, B.T. Miller, A.L. Braswell, P.V. Cupp Jr., and C.S. Hobson. 1999. Conservation status of the southern Appalachian herpetofauna. *Virginia Journal of Science* 50:13-35.

Monarch Joint Venture. Breeding habitat loss. [Consulted 28 May 2015]. Available from <http://www.monarchjointventure.org/threats/breeding-habitat-loss/>

NALCC (North Atlantic Landscape Conservation Cooperative). 2014. *Connecticut River Watershed Landscape Conservation Design Pilot*. U.S. Fish and Wildlife Service fact sheet.

National Geographic. 2010. About one cubic foot. Freshwater, Duck River, Tennessee. Available from <http://ngm.nationalgeographic.com/2010/02/cubic-foot/liittschwager-photography>

NEPARC (Northeast Partners in Amphibian and Reptile Conservation). 2013. Snake fungal disease: frequently asked questions. NEPARC Publication 2013-02. Available from http://www.northeastparc.org/products/pdfs/NEPARC_SnakeFungalDiseaseFAQ.pdf

NFWPCAP (National Fish, Wildlife and Plants Climate Adaptation Partnership). 2012. National fish, wildlife and plants climate adaptation strategy. Washington (DC): Association of Fish and Wildlife agencies, Council on Environmental Quality, Great Lakes Indian Fish and Wildlife Commission, National Oceanic and Atmospheric Administration, and US Fish and Wildlife Service 120 p. Available from <http://www.wildlifeadaptationstrategy.gov/pdf/NFWPCAS-Final.pdf>

NHA (National Hydropower Association). Hydropower for a clean energy future. Available from <http://www.hydro.org/wp-content/uploads/2010/12/NHA-Fact-Sheet-Hydropower-for-a-Clean-Energy-Future.pdf>

Nhancale, B.A. and R.J. Smith. 2011. The influence of planning unit characteristics on the efficiency and spatial pattern of systematic conservation planning assessments. *Biodiversity and Conservation* 20:1821–1835.

Nicholson, C.P., R.D. Tankersley, Jr., J.K. Fiedler, and N.S. Nicholas. 2005. Assessment and prediction of bird and bat mortality at wind energy facilities in the Southeastern United States. Tennessee Valley Authority. Available from http://www.tva.com/environment/bmw_report/bird_bat_mortality.pdf

Niemiller, M.L., B.M. Glorioso, C. Nicholas, J. Phillips, J. Rader, E. Reed, K.L. Sykes, J. Todd, G.R. Wyckoff, E.L. Young, and B.T. Miller. 2006. Status and distribution of the streamside salamander, *Ambystoma barbouri*, in Middle Tennessee. *The American Midland Naturalist* 156 (2):394-399.

Niemiller, M.L. and K.S. Zigler. 2013. Patterns of cave biodiversity and endemism in the Appalachians and Interior Plateau of Tennessee, USA. *PLoS One* 8(5):e64177. Available from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3661478/>

Nislow, K.H., M. Hudy, B.H. Letcher, and E.P. Smith. 2011. Variation in local abundance and species richness of stream fishes in relation to dispersal barriers: implications for management and conservation. *Freshwater Biology* (2011) 56, 2135–2144 doi:10.1111/j.1365-2427.2011.02634.x. Available from http://www.researchgate.net/profile/Benjamin_Letcher/publication/230546030_Variation_in_local_abundance_and_species_richness_of_stream_fishes_in_relation_to_dispersal_barriers_implications_for_management_and_conservation/links/02e7e524f2434a4851000000.pdf

NPS (National Park Service). 2002. Obed Wild and Scenic River Final Climbing Management Plan. Available from <http://www.nps.gov/obed/learn/management/upload/Obed-Final-Climbing-Management-Plan.pdf>

NPS. 2006. Big South Fork National River and Recreation Area/Obed Wild and Scenic River Oil and Gas Management Plan/Environmental Impact Statement. Final Internal Scoping Report, July 2005 (Updated January 2006).

NPS. 2012. Big South Fork National River and Recreation Area and Obed Wild and Scenic River Final Non-Federal Oil and Gas Management Plan, Environmental Impact Statement. [Consulted 4 July 2015a]. OEPC Control Number: FES 12-25. Available from <http://parkplanning.nps.gov/document.cfm?documentID=48597>

NPS. 2014. Big South Fork National River and Recreation Area Mine Drainage EIS Scoping Brochure. [Consulted 30 May 2015]. Available from <http://parkplanning.nps.gov/document.cfm?parkID=354&projectID=42994&documentID=60181>

NPS. 2015a. Great Smoky Mountains, water quality. [Consulted 30 May 2015]. Available from <http://www.nps.gov/grsm/learn/nature/water-quality.htm>

NPS. 2015b. Great Smoky Mountains, air quality. [Consulted 30 May 2015]. Available from <http://www.nps.gov/grsm/learn/nature/air-quality.htm>

NPS. 2015c. Great Smoky Mountains, Hemlock Woolly Adelgid. [Consulted 11 April 2015]. Available from <http://www.nps.gov/grsm/learn/nature/hemlock-woolly-adelgid.htm>

NPS. 2015d. Great Smoky Mountains National Park wildland fire. [Consulted 30 May 2015]. Available from <http://www.nps.gov/grsm/learn/nature/wildlandfire.htm>

Palmer, S.R. 2012. Birds, bats and windmills. The Nature Conservancy. PowerPoint Presentation.

Palmer, S.R. and J.P. Wisby. 2011. Linking conservation priority to wetland and stream mitigation decisions: a watershed planning approach for the Stones River, Tennessee. Environmental Law Institute and The Nature Conservancy 80 p.

Pennsylvania Department of Conservation and Natural Resources. 2015. Forest health factsheet: hemlock woolly adelgid. [Consulted 30 May 2015]. Available from http://www.dcnr.state.pa.us/cs/groups/public/documents/document/dcnr_007179.pdf

Petts, G.E. 2009. Instream flow science for sustainable river management. Journal of the American Water Resources Association 45:1071–1086. doi: 10.1111/j.1752-1688.2009.00360.x. Available from <http://onlinelibrary.wiley.com/doi/10.1111/j.1752-1688.2009.00360.x/abstract>

Powell, J.R. 2003. Response of fish communities to cropland density and natural environmental setting in the Eastern Highland Rim Ecoregion of the Lower Tennessee River Basin, Alabama and Tennessee, 1999. Nashville (TN): USGS Water-Resources Investigations Report 2002-4268. Available from <http://pubs.usgs.gov/wri/wri024268/pdf/wri024268.pdf>

Salafsky, N., R. Margoluis, and K. Redford. 2001. Adaptive management: a tool for conservation practitioners. Washington (D.C.): Biodiversity Support Program.

Salafsky, N., D. Salzer, A.J. Stattersfield, C. Hilton-Taylor, R. Neugarten, S.H.M. Butchart, B. Collen, N. Cox, L.L. Master, S. O'Connor, and D. Wilkie. 2008. A standard lexicon for biodiversity conservation: unified classifications of threats and actions. *Conservation Biology* 22(4):897-911. doi: 10.1111/j.1523-1739.2008.00937.x. Available from <http://cmp-openstandards.org/wp-content/uploads/2014/03/Classification-of-threats-and-actions.pdf>

Schubert, N. 2013. 2013 Golden-winged Warbler monitoring at Hampton Creek Cove State Natural Area, Tennessee. Report submitted to Tennessee Wildlife Resources Agency.

SELC (Southern Environmental Law Center). 2015. Coal mining in the Cumberlands. [Consulted 29 May 2015]. Available from <https://www.southernenvironment.org/cases-and-projects/coal-mining-in-the-cumberlands>

Shute, P.W., R.G. Biggins, and R.S. Butler. 1997. Management and conservation of rare aquatic resources: a historical perspective and recommendations for incorporating ecosystem management. In: Benz, G.W. and D.E. Collins, editors. *Aquatic fauna in peril*, 1st edition. Decatur (GA): Lenz Design & Communications p. 445-466.

Smith, J. 2009. Effects of urbanization on instream habitat and fish assemblages in the Chattanooga metropolitan area, Tennessee-Georgia [thesis]. The University of Tennessee at Chattanooga 129 p. Publication Number 1488604. Available from <http://gradworks.umi.com/14/88/1488604.html>

Smith, R.K., P.L. Freeman, J.V. Higgins, K.S. Wheaton, T.W. FitzHugh, K.J. Ernstom, and A.A. Das. 2002. Priority areas for freshwater conservation action: a biodiversity assessment of the Southeastern United States. The Nature Conservancy. Available from https://www.conservationgateway.org/ConservationPlanning/SettingPriorities/EcoregionalReports/Documents/se_biodiv_assess.pdf

Solis, M.E., C.C. Liu, P. Nam, D.K. Niyogi, J.M. Bandeff, and Y. Huang. 2007. Occurrence of organic chemicals in two rivers inhabited by Ozark hellbenders (*Cryptobranchus alleganiensis bishopi*). *Archives of Environmental Contamination and Toxicology* 53:426-434.

Southeastern Lake Sturgeon Working Group. 2014. Lake Sturgeon Management Plan for the Tennessee and Cumberland Rivers.

Stein, B.A. 2002. States of the Union: ranking America's biodiversity. Arlington (VA): NatureServe. Available from <http://www.natureserve.org/library/stateofunions.pdf>

Stein B.A., P. Glick, N. Edelson, and A. Staudt, editors. 2014. Climate smart conservation: putting adaptation principles into practice. Washington (DC): National Wildlife Federation. Available from http://www.nwf.org/~media/PDFs/Global-Warming/2014/Climate-Smart-Conservation-Final_06-06-2014.pdf

Stein, B.A. and Gravuer, K. 2008. Hidden in Plain Sight: The Role of Plants in the State Wildlife Action Plans. Arlington, Virginia: NatureServe. Available from http://www.natureserve.org/sites/default/files/publications/files/hidden_in_plain_sight_0.pdf

Stein B.A., L.S. Kutner., and J.S. Adams, editors. 2000. Precious heritage: the status of biodiversity in the United States. New York (NY): Oxford University Press 416 p.

Sun, G., S.G. McNulty, J.A.M. Myers, and E.C. Cohen. 2008. Impacts of stresses on water demand and supply across the southeastern United States. Journal of the American Water Resources Association 44(6):1441–1457.

TANSTF (Tennessee Aquatic Nuisance Species Task Force). 2008. Tennessee Aquatic Nuisance Species Management Plan. Nashville (TN): Tennessee Wildlife Resources Agency. Available from <http://www.anstaskforce.gov/State%20Plans/More/Tnplanfinal.pdf>

TDEC (Tennessee Department of Environment and Conservation). 2009. Tennessee 2020, vision for parks, people, and landscapes. Gardner, E.S., J. M. Fly, and S.A. Fritts, editors. Available from <http://www.tn.gov/environment/article/tennessee-2020-plan>

TDEC. 2012. 2012 305(b) Report: the status of water quality in Tennessee. Division of Water Resources, TDEC.

TDF (Tennessee Department of Agriculture, Division of Forestry). 2010. Tennessee forest resource assessment and strategy. Auth. No. 325401. Available from <http://www.tn.gov/agriculture/topic/ag-forests-action-plan>

TDF. Common insect pests of Tennessee Forests. [Consulted 30 May 2015]. Available online at <http://www.tn.gov/agriculture/topic/ag-businesses-plant-pests>

Tennessee Regions' Roundtable Network. 2013. Navigating our future: best practices case studies from the Tennessee Regions' Roundtable Network 52 p. Available from <http://www.cumberlandregiontomorrow.org/wp-content/uploads/2014/02/Navigating-Our-Future.pdf>

The Heinz Center. 2013. Pollinators and the State Wildlife Action Plans: Voluntary Guidance for State Wildlife Agencies. Washington, DC 20 pp. Available from <http://teaming.com/sites/default/files/SWAP%20Pollinator%20Report%20FINAL%204-2013.pdf>

Thurman, L. and B. Terry. 2011. Land use and planning in Tennessee: Part II land use and transportation planning. Nashville (TN): Tennessee Advisory Commission on Intergovernmental Relations 86 p. Available from http://www.state.tn.us/tacir/PDF_FILES/Other_Issues/LandUseAndPlanning.pdf

TNBWG (Tennessee Bat Working Group). 2014. White-nose Syndrome in Tennessee. [Consulted 30 May 2015]. Available from http://www.tnbwg.org/TNBWG_WNS.html

TNC (The Nature Conservancy). 2000. Arnold Air Force Base Invasive Pest Plant Inventory and Management Plan (IMPP). Arnold Air Force Base.

TNC. 2015a. Hemlocks and how to save them, Tennessee. [Consulted 30 May 2015]. Available from <http://www.nature.org/ourinitiatives/regions/northamerica/unitedstates/tennessee/explore/hemlock.xml>

TNC. 2015b. Tennessee Caves. [Consulted July 26, 2015]. Available from <http://www.nature.org/ourinitiatives/regions/northamerica/unitedstates/tennessee/placesweprotect/tennessee-caves.xml>

TN-EPPC (Tennessee Exotic Pest Plan Council). TN-EPPC's cost of invasive exotic plants in Tennessee research results. [Consulted 30 May 2015]. Available from <http://www.tneppc.org>

TNPFC (Tennessee Prescribed Fire Council). [Consulted 4 July 2015]. Available from <http://www.tnpfc.org>

Todd, R.M. 2005. Tennessee alligator gar management plan. Tennessee Wildlife Resources Agency.

TVA (Tennessee Valley Authority). 2006. River Neighbors: ATV users face fines. [Consulted 30 May 2015]. Available from <http://www.tva.com/river/neighbors/nov06/atv.htm>

TVA. Managing river system flows. [Consulted 29 May 2015a]. Available from <http://www.tva.gov/river/lakeinfo/systemwide.htm>

TVA. Water Quality: tailwater improvements, boosting oxygen concentrations. [Consulted 29 May 2015b]. Available from http://www.tva.gov/environment/water/rri_index.htm

TWRA (Tennessee Wildlife Resources Agency). 2000. Alligator Snapping Turtle Restoration Plan. Nashville, TN.

TWRA. 2005. Tennessee's Comprehensive Wildlife Conservation Strategy. Nashville (TN): TWRA. Available from <http://www.state.tn.us/twra/cwcs/tncwcs2005.pdf>

TWRA. 2006. TWRA Nongame and Endangered Species Operational Plan 2006-12. Nashville, TN.

TWRA. 2009. Climate change and potential impacts to wildlife in Tennessee: an update to Tennessee's State Wildlife Action Plan. TWRA Wildlife Technical Report 09-09 106 p. Nashville, TN. Available from <http://www.state.tn.us/twra/pdfs/tncclimatechange.pdf>

TWRA. 2014. The Tennessee Wildlife Resources Agency Strategic Plan 2014-20. Nashville (TN) 81 p. Nashville, TN. Available from <http://www.state.tn.us/twra/pdfs/businessplan.pdf>

TWRA. 2015. Tennessee fishing guide 2015-2016. Knoxville (TN): The Bingham Group 52 p. Available from <http://www.state.tn.us/twra/pdfs/fishguide.pdf>

TWRA, TDEC, and TDA (Tennessee Dept. of Agriculture). 2006. Tennessee Heritage Conservation Trust Fund Act of 2005.

TWRA and USFWS (U.S. Fish & Wildlife Service). 2002. West Tennessee Wildlife Resources Conservation Plan. Nashville, TN.

University of Wisconsin. 2014. Switchgrass production for biomass (Research Brief #51). [Consulted 31 May 2015]. Madison (WI): College of Agricultural and Life Sciences, Center for Integrated Agricultural Systems. Available from <http://www.cias.wisc.edu/switchgrass-production-for-biomass/>

USACE (U.S. Army Corps of Engineers). 2015. Duck River Watershed Assessment. [Consulted 29 May 2015]. Available from <http://www.lrn.usace.army.mil/Media/FactSheets/FactSheetArticleView/tabid/6992/Article/493825/duck-river-watershed-assessment.aspx>

U.S. Census. 2000. Census 2000 Gateway. [Consulted 11 April 2015]. Available from <http://www.census.gov/main/www/cen2000.html>

U.S. Census. 2011. Population distribution and change: 2000 to 2010. By Mackun, P. and S. Wilson. United States Census Bureau: 2010 Census Briefs. Available from <http://www.census.gov/prod/cen2010/briefs/c2010br-01.pdf>

USDA (U.S. Department of Agriculture). 2014. USDA improves forest health by harvesting biomass for energy. [Consulted 5 July 2015]. Available from http://www.fsa.usda.gov/FSA/printapp?fileName=nr_20141216_rel_0270.html&newsType=newsrel

USDOT-FHA (U.S. Department of Transportation, Federal Highway Administration). 2007. Design for fish passage at roadway-stream crossings: synthesis report. Maclean (VA): Federal Highway Administration. Available from <http://www.fhwa.dot.gov/engineering/hydraulics/pubs/07033/07033.pdf>

USFS (USDA Forest Service). 2010. Bioenergy and biobased projects - strategic 2009-2014. Available from <http://www.fs.fed.us/research/docs/priority/bioenergy-strategic-direction.pdf>

USFS. 2013. Forest disturbance processes. [Consulted 30 May 2015]. Available from <http://www.nrs.fs.fed.us/disturbance/pollution/>

USFS. Fire Effects Information System, plant species database. [Consulted 31 May 2015a]. Available from <http://www.fs.fed.us/database/feis/plants/shrub/lesbic/all.html>

USFS. Landscape-scale thresholds of early successional habitat: reconciling biodiversity, public perception, and timber yield in managed forests. [Consulted 28 May 2015b]. Available from <http://www.srs.fs.usda.gov/uplandhardwood/research-topics/duplicates/nifa-grant-loeb.html>

USFS. Southern Research Station Research Areas. [Consulted 28 May 2015c]. Available from http://www.srs.fs.usda.gov/forestops/research_areas/

USFWS (U.S. Fish and Wildlife Service). 2014. White-nose Syndrome: the devastating disease of hibernating bats in North America. Available from https://www.whitenosesyndrome.org/sites/default/files/resource/white-nose_fact_sheet_8-2014_0.pdf

USFWS. 2015. Endangered species permits: NiSource Habitat Conservation Plan. [Consulted 29 May 2015]. Available from <http://www.fws.gov/Midwest/endangered/permits/hcp/nisource/index.html>

USGS (U.S. Geological Survey). 1997. Mercury contamination of aquatic ecosystems. [Consulted 28 May 2015]. Available from http://water.usgs.gov/wid/FS_216-95/FS_216-95.html

USGS. 1997. National water summary on wetland resources. [Consulted 4 July 2015]. Water Supply Paper 2425. Available from https://water.usgs.gov/nwsum/WSP2425/state_highlights_summary.html

USGS. 2013. Environmental flow research in the Tennessee River Basin. [Consulted 28 May 2015]. Available from <http://tn.water.usgs.gov/projects/EcologicalFlows/keyfindings.html>

USGS. 2013. Summary of monitoring and assessments related to environmental flows in USGS Water Science Centers across the U.S. Available from http://water.usgs.gov/coop/enviroflows_summary.pdf

USGS. Effects of urbanization on stream ecosystems. [Consulted 28 May 2015]. Available from <http://water.usgs.gov/nawqa/urban/html/findings/index.html>

- Veni, G., H. DuChene, N.C. Crawford, C.G. Graves, G.N. Huppert, E.H. Kastning, R. Olson, and B.J. Wheeler. 2001. Living with karst: a fragile foundation. Alexandria (VA): American Geological Institute. Available from <http://www.agiweb.org/environment/publications/karst.pdf>
- Walker, G., E. Parish, P. Smith, D. Whitlock, D. Kramar, U. Matthes, and L. Morefield. 2009. Characterization of plant community structure and abiotic conditions on climbed and unclimbed cliff faces in the Obed River Gorge [thesis]. Appalachian State University. Available from <http://www.nps.gov/obed/learn/management/upload/Plant-Cliff-study.doc>
- Warren, M.L. and M.G. Pardew. 1998. Road crossings as barriers to small-stream fish movement. Transactions of the American Fisheries Society 127:637-644. Available from <http://www.stream.fs.fed.us/fishxing/fplibrary/Warren%20and%20Pardew-1998.pdf>
- Webbers, A. 2003. Public water-supply systems and associated water use in Tennessee, 2000. Nashville (TN): U.S. Geological Survey Water-Resources Investigations Report 03-4264. Available from <http://pubs.usgs.gov/wri/wri034264/>
- Williams, C.E., R.D. Bivens, B.D. Carter, and C.E. Skelton. 2014. Checklist of Tennessee crayfishes (Decapoda: Cambaridae). Morristown (TN): Tennessee Wildlife Resources Agency, in-house report.
- Wisby, J.P. and S.R. Palmer. 2015. Database Development and Spatial Analyses in Support of Tennessee's State Wildlife Action Plan. The Nature Conservancy, Nashville, TN.
- Woody, L. 2011. Back to the Future. Tennessee Wildlife Magazine.
- Yarbrough, J. 2013. TVA's Aquatic Resources Stewardship & "Vital Signs" Monitoring. Tennessee Valley Authority.
- Young B.E., E. Byers, K. Gravuer, K. Hall, G. Hammerson, A. Redder, J. Cordeiro, and K. Szabo. 2011. Guidelines for using the NatureServe Climate Change Vulnerability Index, version 2.1. Arlington (VA): NatureServe.
- Young, B.E., K.R. Hall, E. Byers, K. Gravuer, G. Hammerson, A. Redder, and K. Szabo. 2012. Rapid assessment of plant and animal vulnerability to climate change. In: Brodie, J., E. Post and D. Doak, editors. **Wildlife Conservation in a Changing Climate**. Chicago (IL): University of Chicago Press. p. 129-152.

Glossary of Terms

acid deposition - also known as acid rain, is an environmental phenomenon caused by air pollutants, mainly nitrates (NOX) and sulfates (SOX) primarily from burning fossil fuels. Acid deposition is characterized by extremely high acidity which is particularly harmful to sensitive ecosystems.

aquatic ecological system - dynamic spatial assemblages of ecological communities (e.g. rivers, streams, and lakes) with similar geomorphological patterns tied together by ecological processes (e.g. hydrologic and nutrient regimes, access to floodplains) or environmental gradients (e.g. temperature, chemical and habitat volume), and form a cohesive, distinguishable unit on a hydrography map.

barrens - Tennessee's barrens ecosystems are typically mosaics of open canopy woodlands, or lacking trees altogether, with a dominance of perennial grasses and herbaceous species

bioaccumulation - the process by which organisms can take up contaminants more rapidly than their bodies can eliminate them.

biodiversity - the full range of natural variety and variability within and among living organisms, and the ecological and environmental complexes in which they occur. It encompasses multiple levels of organization, including genes, species, communities, and ecological systems or ecosystems.

bioenergy - a general term that encompasses both biofuel and biomass as sources of energy generation.

biofuels - biofuels, generally defined as liquid fuels derived from biological materials, can be made from plants, vegetable oils, forest products, or waste materials. The raw materials can be grown specifically for fuel purposes, or can be the residues or wastes of existing supply and consumption chains, such as agricultural residues or municipal garbage.

biomagnification - also known as bioamplification or biological magnification, is the sequence of processes in an ecosystem by which higher concentrations of a particular chemical, such as DDT or mercury, are reached in organisms higher up the food chain, generally through a series of prey-predator relationships.

biomass - solid organic materials produced in a renewable manner for energy generation. Two categories of biomass fuels, woody fuels and animal wastes, comprise the vast majority of available biomass fuels.

buffer - conservation buffers are small areas or strips of land in permanent vegetation, designed to slow water runoff, provide shelter and stabilize riparian or other habitat areas. Strategically placed buffer strips in the agricultural landscape can effectively mitigate the movement of sediment, nutrients, and pesticides within farm fields and from farm fields.

conservation action - any act taken to directly abate a stress or source of stress to a target species or habitat, or to prevent the future development of a stress upon a species or its habitat.

conservation goal - in conservation planning, the number and spatial distribution of on-the-ground occurrences of targeted species, natural communities, and ecological systems that are needed to adequately conserve the target in an ecoregion.

Conservation Opportunity Area - Priority areas for conservation, spatially depicted on the landscape, that offer the best opportunities and potential for conservation of species of greatest conservation need and their habitats.

dam - any structure that impounds water.

decline/declining - the historical or recent decrease of a conservation target through all or part of its range. Declining species exhibit significant, long-term decreases in habitat and/or numbers, are subject to a high degree of threat, or may have unique habitat or behavioral requirements that expose them to great risk.

diadromous - Fish that can live in and are migratory between fresh waters and salt waters.

disjunct - distributional range of a species or community which is found in an ecoregion a significant distance from its primary range in other disconnected ecoregions. Disjunct species have populations that are geographically isolated.

distribution pattern - the overall pattern of occurrence for a particular conservation target. In ecoregional planning, distribution patterns are often described in terms of the relative proportion of the target's natural range occurring within a given ecoregion (i.e. endemic, limited, disjunct, peripheral, and widespread).

dry bed retention dam - a dry detention basin temporarily detains and stores collected stormwater runoff for a period of time, releasing the stormwater slowly to reduce flooding and remove pollutants. It is referred to as "dry" because it dries out between rain events. Pollutants are removed by allowing particulates and solids to settle out from the water. Detention does not normally occur until the inflow rate exceeds the design outflow rate.

dry-mesic (forest) - Oak or oak-hickory forest type (typically), characterized by well-drained soils and episodic fire.

ecological systems - ecological systems are dynamic assemblages of native plant and/or animal communities that 1) occur together on the landscape or in the water, 2) are tied together by similar ecological processes (e.g., fire, hydrology), underlying environmental features (e.g., soils, geology), or environmental gradients (e.g., elevation).

ecoregion - a relatively large geographic unit of land and water defined by the climate, vegetation, geology, and other ecological and environmental patterns.

element occurrence (EO) - a term originating from methodology of the Natural Heritage Program network that refers to species, natural communities, or other entities (e.g. migratory bird stopovers, ecological systems) of biodiversity that serve as both conservation targets and as units for organizing and tracking information.

endemic - distributional range of a species or community which primarily or only occurs in one specific area

eutrophication - the process by which a body of water acquires a high concentration of nutrients, especially phosphates and nitrates. These typically promote excessive growth of algae. As the algae die and decompose, high levels of organic matter and the decomposing organisms deplete the water of available oxygen, causing the death of other organisms, such as fish. Eutrophication may be a natural, slow-aging process for a water body, but human activity can greatly speed up or amplify the process.

global rank - a numeric assessment of a biological element's relative imperilment and conservation status across its range of distribution ranging from G1 (critically imperiled) to G5 (secure). Assigned by the Natural Heritage Programs, global ranks for species and natural communities are determined primarily by the number of occurrences or total area of coverage (communities only), modified by other factors such as condition, historic trend in distribution or condition, vulnerability, and threats.

Greatest Conservation Need - a designation determined by each state's fish and wildlife agency, which identifies the species and their associated habitats that are most at risk, threatened, or deserving of conservation action for other reasons.

green infrastructure - a strategically planned and managed network of natural lands, working landscapes, and other open spaces that conserve ecosystem values and functions and provide associated benefits to human populations.

habitat fragmentation - occurs when large ecosystems are separated into smaller pieces, separating and isolating groups of species and habitats from one another, which impacts breeding patterns and may lead to inbreeding.

headwaters - the source area where a river or stream begins in a landscape.

hibernacula - (single hibernaculum) location where a creature seeks shelter over the winter, such as bats using a cave.

incompatible management practices - incompatible practices modify habitat composition, type, and/or ecological processes in a way that is not compatible with the needs of target wildlife or plant species.

instream flow - the natural variations of water levels in rivers and streams, critical to the maintenance of biodiversity and other values, such as recreation, in aquatic ecosystems.

invasive exotic species - nonindigenous species which have been introduced either intentionally or accidentally into areas outside their natural range and that have the capacity to outcompete native species either reproductively or for natural resources.

karst - landscape formed from the dissolution of soluble rocks including limestone, dolomite and gypsum. Karst is characterized by sinkholes, caves, and underground drainage systems.

keystone species - species with an important role in ecosystem function, such as top carnivores.

limited - distributional range of a species or natural community which occurs in the ecoregion and within a few other adjacent ecoregions.

mesophytic (forest) - a forest that generally receives a moderate amount of moisture.

natural communities - terrestrial plant communities of definite floristic composition, uniform habitat conditions, and uniform physiognomy. Natural communities are defined by the finest level of classification, the “plant association” level of the National Vegetation Classification. Like ecological systems, natural plant communities are characterized by both a biotic and abiotic component.

natural process - processes in nature that play a vital role in ecosystem function by causing change or disturbance on a cyclical basis, such as flooding, fire, or plant succession. Habitat restoration can sometimes be achieved by incorporating natural processes into conservation designs.

occurrence - a spatially referenced location of a species or a location of a natural plant community or ecological system. Many occurrences are tracked by the various Natural Heritage Programs and are known as Element Occurrences. Occurrences may also be more loosely defined locations delineated through the definition/mapping or other spatial data or through the identification of areas by experts.

parcelization - generally refers to the division of ownerships that result in smaller holdings of land which, in turn, results in constrained management options and potentially adverse effects on ecosystem health and wildlife habitat. Parcelization is caused by subdividing large tracts into smaller forest tracts, ranchettes for residential use, or sale of large tracts to buyers who further subdivide the land.

prescribed fire - fire that is purposely lit under controlled conditions and guided by a burn plan to improve ecological health, reduce fuel loads that contribute to catastrophic fire, or achieve desired habitat conditions.

propagation - the breeding of plants or animals for reintroduction into suitable habitats as supplementation of existing populations or as a means of restoring locally extirpated populations.

problem - the combined concept of ecological stresses to a target and the sources of that stress to the target.

refugia - (single refugium) areas in which a population of organisms can survive through a period of unfavorable conditions.

rockhouse - a rockhouse is an area that typically has been eroded away by a stream under a large rock or bluff line. Common on the Cumberland Plateau, they are typically found in sandstone where the old stream meandered next to a rock bluff. .

sinkhole - a cavity in the ground, especially in limestone bedrock, caused by water erosion and providing a route for surface water to disappear underground.

source (of stress) - an extraneous factor, either human (i.e. activities, policies, land uses) or biological (e.g. non-native species), that infringes upon a conservation target in a way that results in stress.

species richness - the number of different species represented in an ecological community, landscape or region.

stress - something which impairs or degrades the size, condition, or landscape context of a conservation target, resulting in reduced viability.

split estate - in split estate situations, the surface rights and subsurface rights (such as the rights to develop minerals) for a piece of land are owned by different parties.

sprawl - low-density decentralized development at the fringes of a central city. It is characterized by a spreading out of development over a wide area with little or no connectivity to the contiguously developed area.

stream connectivity - streams, from headwaters to large rivers, provide pathways for the transport of a wide range of materials, energy and aquatic life. This transport moves upstream and downstream, between surface and groundwater, between channel and floodplain and in time, i.e. seasonal flooding or scouring.

stocking - relative to fish or other aquatic organisms, the practice of raising organisms in a hatchery and releasing them into appropriate lake, stream, or other aquatic habitat.

substrate embeddedness - the degree to which fine sediments surround coarse substrates on the surface of a streambed.

succession - the process of change in the species structure of an ecological community over time. Generally resulting in increasing structural complexity over time until a community becomes relatively stable or self-perpetuating in the absence of disturbance.

tailwater - waters located immediately downstream from a hydraulic structure, most often a dam.

target - specific components of biodiversity which serve as the focus of conservation, including the development and prioritization of conservation strategies. Conservation targets consist of ecological systems, natural communities, and species.

translocation - the capture, transport, and release or introduction of species, habitats or other ecological material from one location to another.

troglobite - an animal that cannot survive outside its cave environment.

troglophile - an animal able to live its entire life in a cave, but usually maintain some of their senses such as partial pigmentation and are usually only partially blind.

trogloxene - an animal that uses caves for shelter but does not complete its life cycle in them, for example bats.

umbrella species - a species or group of species, such as forest interior dwelling birds, whose habitat needs overlap those of other animals and plants.

viable/viability - the ability of a species to persist for many generations, or a natural community or ecological system to persist over some time period. An assessment of viability will often focus on the minimum area and number of occurrences and presence of natural processes necessary for persistence.

viewshed - the geographical area that is visible from a location. It includes all surrounding points that are in line-of-sight with that location and excludes points that are beyond the horizon or obstructed by terrain and other features (e.g., buildings, trees).

volant - able to fly or glide.

widespread - distributional range of a species or natural community which is typically found in the ecoregion, but common in many others also; the bulk of distribution may be elsewhere however.

vulnerability - the inability to withstand the effects of a hostile environment, generally measured by degree of exposure to hostile elements or change, inherent sensitivity to that exposure, and adaptive capacity to accommodate change.



Credits clockwise from top left: **Lamprey on a Striped Shiner** - Jeremy Monroe, **Fisheries Illustrated**; Streamside Salamander - Matthew Niemiller; Western Sandpiper - Chris Sloan; **Bog Turtle** - Jeffrey Basinger, **Fisheries Illustrated**; Upland Burrowing Crayfish - Carl Williams, TWRA; Little Brown Bats - Chris Ogle, TWRA